

## Geology

### Rare earth element chemistry of zircon and its use as a provenance indicator

Paul W.O. Hoskin and Trevor R. Ireland

*Geology* 2000;28:627-630

doi: 10.1130/0091-7613(2000)28<627:REECOZ>2.0.CO;2

---

**Email alerting services**

click [www.gsapubs.org/cgi/alerts](http://www.gsapubs.org/cgi/alerts) to receive free e-mail alerts when new articles cite this article

**Subscribe**

click [www.gsapubs.org/subscriptions/](http://www.gsapubs.org/subscriptions/) to subscribe to *Geology*

**Permission request**

click <http://www.geosociety.org/pubs/copyrt.htm#gsa> to contact GSA

Copyright not claimed on content prepared wholly by U.S. government employees within scope of their employment. Individual scientists are hereby granted permission, without fees or further requests to GSA, to use a single figure, a single table, and/or a brief paragraph of text in subsequent works and to make unlimited copies of items in GSA's journals for noncommercial use in classrooms to further education and science. This file may not be posted to any Web site, but authors may post the abstracts only of their articles on their own or their organization's Web site providing the posting includes a reference to the article's full citation. GSA provides this and other forums for the presentation of diverse opinions and positions by scientists worldwide, regardless of their race, citizenship, gender, religion, or political viewpoint. Opinions presented in this publication do not reflect official positions of the Society.

---

**Notes**

# Rare earth element chemistry of zircon and its use as a provenance indicator

Paul W.O. Hoskin\* Research School of Earth Sciences, Australian National University, Canberra, ACT 0200, Australia  
 Trevor R. Ireland\* Department of Geological and Environmental Sciences, Stanford University, Stanford, California 94305-2115, USA

## ABSTRACT

Sedimentary mineral assemblages commonly contain detrital zircon crystals as part of the heavy-mineral fraction. Age spectra determined by U-Pb isotopic analysis of single zircon crystals within a sample may directly image the age composition—but not the chemical composition—of the source region. Rare earth element (REE) abundances have been measured for zircons from a range of common crustal igneous rock types from different tectonic environments, as well as kimberlite, carbonatite, and high-grade metamorphic rocks, to assess the potential of using zircon REE characteristics to infer the rock types present in sediment source regions. Except for zircon with probable mantle affinities, zircon REE abundances and normalized patterns show little intersample and intrasample variation. To evaluate the actual variation in detrital zircon REE composition in a true sediment of known mixed provenance, zircons from a sandstone sample from the Staffjord Formation (North Sea) were analyzed. Despite a provenance including high-grade metasediment and granitoids and a range in zircon age of 2.82 b.y., the zircon REEs exhibit a narrow abundance range with no systematic differences in pattern shape. These evidences show zircon REE patterns and abundances are generally not useful as indicators of provenance.

**Keywords:** provenance, rare earths, trace elements, zircon.

## INTRODUCTION

Sediments and sedimentary rocks, in recording past erosion and deposition, provide a record of tectonic activity. Zircon is an often studied component of detrital assemblages because it may withstand mechanical and chemical breakdown over billions of years. Another important characteristic is that zircon crystallizes with an extremely high U/Pb ratio and retains the daughter products of U and Th radioactive decay so that the array and frequency of U-Th-Pb ages measured on detrital zircon populations yields information about the ages of crustal elements in the source region and the clastic pathway (Roback and Walker, 1995; Machado et al., 1996; Ireland et al., 1998; Sircombe, 1999).

A potentially powerful complement to the chronology and typology (Dabard et al., 1996) of detrital zircon would be the identification of source-rock lithology by using zircon trace element composition. The populations of zircon crystals within a sediment or sedimentary rock could then potentially be used to determine both the ages and rock types in the source region. This dual approach of assessing zircon isotope and trace element chemistry would have enormous potential in determining sediment provenance, and its applications to studies of, for example, uplift and erosion, tectonics, and magma genesis.

\*E-mail: Hoskin—Papanui@xtra.co.nz; Ireland—tri@pangea.stanford.edu.

## BACKGROUND PRINCIPLES AND CONSIDERATIONS

Any mineral phase will indicate its whole-rock source if (1) the mineral composition can be related to a melt composition by the appropriate use of mineral/melt partition coefficients ( $K_d$  values); (2) the mineral preserves elemental ratios that are also retained in the whole rock; or (3) chemical features of a mineral can be distinguished and systematically related to the rock type of origin.

Successful application of  $K_d$  values to igneous minerals for the purpose of provenance tracing relies upon knowing the  $K_d$  values under a range of magmatic conditions, accurate measurement of the elemental abundances, and the validity of assumptions that (1) the mineral was saturated in the melt at the liquidus and (2) equilibrium mineral/melt partitioning was attained. Saturation calculations for zircon (Watson and Harrison, 1983) and modeling of cogenetic rock suites reveal that zircon is only likely to be a liquidus phase in felsic melts and in some intermediate igneous compositions. Zircon found in mafic and many intermediate rocks probably grew in trapped melt pools, far from the liquidus. The assumption of equilibrium element trace partitioning between zircon and melt in most cases is probably invalid because surface enrichment and growth entrapment (i.e., disequilibrium partitioning) of the rare earth elements (REEs, including Y) and other elements is inevitable (Watson, 1996). Moreover, recent experimental studies reveal that REE partitioning into zircon is not a simple func-

tion of REE<sup>3+</sup> ionic radii, but is greatly affected by structural strain by the REE as well as charge-balancing elements at both the Zr and Si lattice sites (Hanchar et al., 2000; Finch et al., 2000).

Using a different approach (that of 3 in the previous list), Heaman et al. (1990) showed that the trace element abundances of zircon from different rock types may cluster into discrete groupings on Harker-style discrimination plots. However, for the five rock groupings of Heaman et al. (1990) there is no clear discrimination on the basis of Lu, Sc, Th/U, or Lu/Sm alone, and only partial discrimination based on reported Hf abundances. Other previous and subsequently published zircon trace element analyses do not conform to the chemical discriminants of Heaman et al. (1990) (e.g., Murali et al., 1983; Hinton and Upton, 1991; Ireland and Wlotzka, 1992; Barbey et al., 1995; Belousova et al., 1998; Hoskin, 1998). Heaman et al. (1990) conceded that if zircon is to become widely used as a tracer, studies must be made of changes in zircon composition as a function of changes in lithology and degree of fractional crystallization.


Here we present microprobe SIMS (secondary ion mass spectrometry) and laser ablation ICP-MS (inductively coupled plasma–mass spectrometry) analyses of well-characterized zircon populations to assess the efficacy of zircon REE abundances and patterns as a provenance indicator. Raw data were normalized against NIST SRM 610. Reproducibility of elemental abundances was monitored by analysis of Mud Tank carbonatite and Boggy Plain zoned pluton zircon by both analytical techniques. Full analytical details are given elsewhere.<sup>1</sup> The results are tested by combined in situ U-Pb isotope and REE analysis of single detrital zircons from a petroleum-well sandstone sample from the Brent Field, northern North Sea.

## RESULTS

To assess changes to the REE abundances and patterns of zircon that crystallized within a magma undergoing fractional crystallization, zircon populations from six whole-rock samples

<sup>1</sup>GSA Data Repository item 200068, Table 1, Analysis techniques and data (ppm) for analyzed zircon in Figure 1, is available on request from Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301-9140, editing@geosociety.org, or at [www.geosociety.org/pubs/ft2000.htm](http://www.geosociety.org/pubs/ft2000.htm).

Data Repository item 200068 contains additional material related to this article.



**Figure 1.** Chondrite-normalized rare earth element patterns for zircon from following sources. **A:** Boggy Plain zoned pluton, Australia. **B:** Blind Gabbro, Australia, and igneous charnockite from Mawson Coast, Antarctica. **C:** Gabbroic ophiolite, Syros, Greece, and plagiogranite, Manilla, Australia. **D:** Kimberlite from Jwaneng, Botswana; carbonatite from Phalaborwa Complex, South Africa; and carbonatite from Mud Tank carbonatite complex, Australia. **E:** High-grade metamorphic rocks from Sri Lanka; north Queensland, Australia; the Vosges Mountains, France (Schaltegger et al., 1999). **F:** Sandstone from Statfjord Formation, North Sea. Chondrite values are from McDonough and Sun (1995).

from the Boggy Plain zoned pluton, Australia, were analyzed. The pluton is concentrically zoned with olivine-bearing diorite on the margin, ranging through granodiorite, adamellite, to aplite in the center (Wyborn et al., 1987). Whole-rock samples range in composition from 52.24 wt% (diorite) to 74.80 wt%  $\text{SiO}_2$  (aplite). Zircon occurs in all compositional zones of the pluton (Hoskin et al., 2000). An expectation of zircon chemistry within this closed, fractionating magma system is that it will reflect compositional changes within the bulk magma as differentiation proceeds, in particular that early crystallizing zircon will be enriched in elements that have bulk distribution coefficients,  $D$ , <1, such as the REEs in this system. Although this is the case for apatite within the pluton (Hoskin et al., 2000), the zircon REE patterns do not vary systematically, and the abundance range measured for a single population completely or significantly overlaps the range for all other popula-

tions. This result is illustrated in Figure 1A, in which chondrite-normalized REE patterns for zircon from the diorite and aplite zones are plotted; sets of zircon REE patterns from four samples from the granodiorite and adamellite zones fall within the same abundance range and are omitted for clarity.

Accessory-mineral assemblages in mafic igneous rocks may include zircon. A 2 kg sample of the Blind Gabbro, Australia, yielded in excess of 1000, 50–250- $\mu\text{m}$ -long subhedral zircon crystals. The zircon REE pattern (Fig. 1B) has a shape similar to that for zircon from the Boggy Plain zoned pluton, but abundance levels of the light REEs (La–Nd) and middle REEs (Sm–Tb) are slightly higher. In contrast to Blind Gabbro, which may have formed above a subduction zone (Miller and Gray, 1996), the mafic igneous charnockite of Ufs Island, East Antarctica, represents an orogenic magma that probably formed by melting of thickened crust (Young and Black,

1991). Zircons in this rock exhibit REE patterns that parallel the range defined by Blind Gabbro zircon (Fig. 1B), but have lower abundances that largely overlap with zircon from the Boggy Plain zoned pluton.

The heavy REE (Dy–Lu) patterns of zircon from Blind Gabbro and Mawson igneous charnockite have a slight concave-down curvature. This is exhibited also for igneous zircon from a metamorphosed gabbroic ophiolite (Fig. 1C) on the island of Syros, Greece (Baldwin, 1996). There is considerable overlap of REE abundances for zircon from the ophiolite sample with zircon from an ophiolitic plagiogranite sample (from Manilla, Australia; Aitchison and Ireland, 1995), which does not have heavy REE curvature, suggesting that curvature in zircon heavy REE patterns from mafic rocks (Figs. 1B and 1C) may be due to heavy REE depletion of melt by crystallization of phases such as clinopyroxene and orthopyroxene. Plagiogranite zircons in this study have small Eu anomalies ( $\text{Eu}/\text{Eu}^*$  averages 0.90).

Samples from kimberlite and carbonatite are included in this survey (Fig. 1D), although detritus from such rocks will be minor constituents in most sediments, and most probably absent. Large zircon crystals (to 1 cm) from kimberlite at Jwaneng, Botswana, are heavy REE enriched but much less so than for other igneous zircons (Figs. 1A–1C). The heavy REE pattern is approximately flat ( $[\text{Yb}/\text{Gd}]_N = 0.8\text{--}1.4$ ). These REE characteristics are observed for other African kimberlitic zircons (Belousova et al., 1998), and the near-flat heavy REE patterns are characteristic of zircon from the mantle-derived MARID (mica-amphibole-rutile-ilmenite-diopside) suite ( $[\text{Yb}/\text{Gd}]_N = 1.3\text{--}3.7$ ; Hoskin, 1998). Zircon from the Mud Tank carbonatite complex, Australia, occurs as megacrysts, sometimes >30 cm long. Fragments of a centimeter-sized zircon crystal were analyzed by Hanchar and Hoskin (1998) and have an REE pattern similar to that of Jwaneng kimberlite zircon, although the heavy REE pattern is steeper ( $[\text{Yb}/\text{Gd}]_N = 7.2\text{--}9.7$ ; Fig. 1D). Neither zircon from the Jwaneng kimberlite or the Mud Tank carbonatite have significant Eu anomalies ( $\text{Eu}/\text{Eu}^* > 0.59$ ). Although the Mud Tank carbonatite was emplaced at mid-crustal levels, the zircon REE pattern may indicate mantle affinities or reflect crystal growth under extremely high water and fluorine fugacities (Currie et al., 1992). We separated 10 small (~30  $\mu\text{m}$ ), anhedral zircon crystals from a slab of carbonatite from the central carbonatite-bearing area of the Phalaborwa Complex, South Africa. The zircons contain numerous opaque inclusions, and only one good analysis was obtained (Fig. 1D). The REE characteristics of this zircon resemble REE patterns from silicic whole-rock samples and contrast with zircon REE patterns from the Mud Tank carbonatite. It is possible that the Phal-

aborwa zircons formed in silica-rich selvages within the carbonatite magma or were derived from silicate magma that is believed to have mixed with the carbonatite (Eriksson, 1989).

Zircon can form by a variety of different processes in rocks undergoing metamorphism, for example, by net-transfer reactions, or solid-state recrystallization, or in low-volume anatectic melts. Schaltegger et al. (1999) documented zircon grown in the presence of aqueous fluids and anatectic melts in high-grade metamorphic rocks from the Vosges Mountains, eastern France. Despite some variation in zircon REE patterns for different rock samples, they found that zircon formed by these different processes has near-identical REE abundances and patterns (Schaltegger et al., 1999, their Fig. 7). The typical REE pattern and range measured by these workers are shown in Figure 1E. Superimposed are REE patterns measured for metamorphic zircon SL1 from the gem gravels of Sri Lanka and metamorphic zircon from granulites of the Georgetown region, north Queensland, Australia. The north Queensland zircons are characterized by nearly flat light REE patterns ( $[Sm/La]_N = 1.3-8.2$ ) and atypical steepness from Gd to Er ( $[Er/Gd]_N = 11-34$ ) relative to other zircon REE patterns. These characteristics are related to the process of formation by solid-state recrystallization of protolith zircon. Partial solid-state recrystallization has been shown to produce flatter light REE patterns and steep heavy REE patterns, and to leave a "memory" of the preexisting chemistry and zoning of the protolith zircons (Hoskin and Black, 2000). The REE patterns of metamorphic zircon from Sri Lanka and France overlap in pattern shape and abundance range and are generally indistinguishable from zircon from silicic crustal igneous rocks (Figs. 1A-1C).

To evaluate the actual variation in detrital zircon REE composition in a real sediment, zircons from a sandstone from the North Sea were analyzed. The sandstone is from the Statfjord Formation (Zone A) and was sampled at 3011 m downhole in Brent Field well 211/29-6 located on the seafloor between Scotland and Norway. Evidence from Sm-Nd isotopes, paleocurrent data, heavy-mineral assemblages, detrital-garnet composition, and detrital-zircon U-Pb isotope spectra indicate that the source region of this sample comprised high-grade metasediment and Caledonian granites (Morton, 1992; Dalland et al., 1995; Morton and Berge, 1995; Morton et al., 1996). The REE abundances for concordant detrital zircon from each of the three main age groups identified by Morton et al. (1996) were analyzed (Fig. 1F). Despite different crustal elements in the source region and an age range of 2.82 b.y. for the zircons we analyzed, there are no systematic differences for zircon REE chemistry within or between the age groupings. This result cannot be attributed to a chemically homogeneous source region or derivation of detritus from

a single source, but confirms the apparent monotony of REE patterns and abundances in zircon derived from a range of common crustal rock types (Figs. 1A-1C and 1E).

Other trace elements were also measured for zircons in Figure 1. The Zr/Hf ratio is highest for zircon from Jwaneng kimberlite (Zr/Hf = 75-168) and Phalaborwa carbonatite (Zr/Hf = 253). All other zircon populations have Zr/Hf values that range across and near the chondritic value of ~37. Apart from high Zr/Hf values in kimberlite and carbonatite, there are no systematic differences between other zircon populations according to source-rock composition. Phosphorus values range considerably; the lowest abundances were measured for zircon from Mawson igneous charnockite (31-77 ppm), Manilla plagiogranite (24-59 ppm), and Mud Tank carbonatite (55-61 ppm). These values are well within the 20-110 ppm range that Belousova et al. (1998) considered to be distinctive for kimberlitic zircons.

Geochronologists have recognized for some time that there is variation in the zircon Th/U elemental ratio, and have used this as a first-order discriminant between igneous and metamorphic zircon (e.g., Ahrens et al., 1967; Williams et al., 1996). Metamorphic zircons in this study (Fig. 1E) have Th/U ratios of 0.01-0.08, whereas zircons of igneous origin (Figs. 1A-1D) range from 0.16 to 2.37. Values greater than Th/U = 1 are uncommon for most igneous zircons; typical values are within the range 0.4-1.0. Due to order of magnitude range in Th/U ratio often found in single zircon populations, we consider the Th/U ratio to be of limited use as a discriminant.

### CONTROLS ON ZIRCON REE CHARACTERISTICS

Although recognizing minor differences between some zircon populations, we have demonstrated that the REE chemistries of zircon from a wide range of crustal rock types and tectonic settings exhibit remarkable similarity. From a North Sea sediment sample of known provenance, we have shown that zircon REE patterns and abundances are not generally useful as indicators of provenance.

A similar conclusion was reached by Maas et al. (1992) in their attempt to determine the provenance of Earth's oldest-known zircons, and by Sawka (1988), who documented zircon REE uniformity across the McMurray Meadows pluton of the Sierra Nevada, United States. There are at least two possible explanations for the apparent monotony of zircon REE characteristics. First, the composition of most silicate melts at the point of zircon saturation can be considered as broadly "granitic." This is true even for trapped intercumulus melt in mafic granitoids, which will evolve toward a granitic composition and perhaps zircon saturation by in situ fractionation. It may be that the REE chemistries of these granitic melts are sufficiently similar to impart REE char-

acteristics to crystallizing zircon that are not significantly different from zircons from a sample with different whole-rock chemistries. However, this is unlikely because the REE chemistry of granitic melts differs depending upon a range of factors, including ambient intensive parameters and crystallization history.

A second explanation for the similarity of zircon REE abundances may relate to lattice strain and charge-balance requirements for REE substitution into the zircon lattice. Incorporation of REEs into the zircon structure is thought to be dominated by the xenotime coupled substitution mechanism (Speer, 1982):  $(Y + REE)^{3+} + P^{5+} = Zr^{4+} + Si^{4+}$ , although this mechanism has now been shown to have strain-related limits (Hanchar et al., 2000; Finch et al., 2000). A limit on the amount of light REE<sup>3+</sup> substitution for Zr<sup>4+</sup> is due to strain at the Zr site and may partly explain (in addition to ionic radii) why La-Nd abundances for all zircon in Figure 1 typically range between 10<sup>-1</sup> and 10<sup>1</sup> times chondrite abundance despite many having grown in melt with strong light REE enrichment (e.g., Jwaneng kimberlite and Mud Tank carbonatite). The heavy REEs have smaller ionic radii that more closely match that of Zr<sup>4+</sup>; incorporation of these elements in zircon is potentially limited by strain at the Si site by incorporation of charge compensating P<sup>5+</sup>. This limit is not reached for most zircon populations in Figure 1 because the mole percent (mol%) of REE exceeds that of P (the xenotime substitution mechanism requires the mol% ratio of REE to P to be unity). This observation suggests that other substitution mechanisms are operating to charge balance the REE, probably in addition to the xenotime mechanism. For zircon from the Boggy Plain zoned pluton, xenotime-type mechanisms involving minor abundances of interstitial cations are believed to provide charge balance for REE in excess of P (Hoskin et al., 2000). Other mechanisms may involve coupled substitution of Nb, Ta, and the REEs at the Zr site, although abundances of Nb and Ta in zircon are typically very low (<10 ppm).

It is possible that heavy REE abundances for most zircon populations in Figure 1 are retarded by the lack of sufficient P<sup>5+</sup> substitution into the mineral lattice, but that further heavy REE substitution is possible by other mechanisms, yet these also are limited by lattice strain. The combined effect of these restricted substitution mechanisms for REE incorporation in zircon is the imposition of REE characteristics that are similar for populations derived from different whole-rock sources.

### CONCLUSIONS

Results from this study demonstrate that the REE chemistry of zircons from a wide range of crustal rock types and tectonic settings exhibit remarkable similarity, and therefore zircon chemistry is not generally useful as an indicator of sedi-

ment or inherited zircon provenance. An exception to this conclusion is zircon with mantle affinities (kimberlite, MARID suite, and perhaps the Mud Tank carbonatite), which have REE:P mol% ratios of 1:1, lower REE abundances, and flatter normalized patterns compared to crustal zircon populations. However, in addition to age data, important provenance information might be obtained by investigation of the occurrence and chemistry of inclusions hosted by detrital zircon.

#### REFERENCES CITED

- Ahrens, L.H., Cherry, R.D., and Erlank, A.J., 1967, Observations on the Th-U relationship in zircons from granitic rocks and from kimberlites: *Geochimica et Cosmochimica Acta*, v. 29, p. 711–716.
- Aitchison, J.C., and Ireland, T.R., 1995, Age profile of ophiolitic rocks across the late Palaeozoic New England orogen, New South Wales: Implications for tectonic models: *Australian Journal of Earth Sciences*, v. 42, p. 11–23.
- Baldwin, S.L., 1996, Contrasting *P-T-t* histories for blueschists from the western Baja terrane and the Aegean: Effects of synsubduction exhumation and backarc extension, in *Bebout, G.E., et al., eds., Subduction: Top to bottom: American Geophysical Union Geophysical Monograph 96*, p. 135–141.
- Barbey, P., Allé, P., Brouand, M., and Albarède, F., 1995, Rare-earth patterns in zircons from the Manaslu granite and Tibetan slab migmatites (Himalaya): Insights in the origin and evolution of a crustally derived granite magma: *Chemical Geology*, v. 125, p. 1–17.
- Belousova, E.A., Griffin, W.L., and Pearson, N.J., 1998, Trace element composition and cathodoluminescence properties of southern African kimberlitic zircons: *Mineralogical Magazine*, v. 62, p. 355–366.
- Currie, K.L., Knutson, J., and Temby, P.A., 1992, The Mud Tank carbonatite complex, central Australia—An example of metasomatism at mid-crustal levels: *Contributions to Mineralogy and Petrology*, v. 109, p. 326–339.
- Dabard, M.P., Loi, A., and Peucat, J.J., 1996, Zircon typology combined with Sm-Nd whole-rock isotope analysis to study Brioverian sediments from the Armorican Massif: *Sedimentary Geology*, v. 101, p. 243–260.
- Dalland, A., Mearns, E.W., and McBride, J.J., 1995, The application of samarium-neodymium (Sm-Nd) provenance ages to correlation of biostratigraphically barren strata: A case study of the Statfjord Formation in the Gullfaks Oilfield, Norwegian North Sea, in *Dunay, R.E., and Hailwood, E., eds., Dating and correlating biostratigraphically-barren strata: Geological Society [London] Special Publication 89*, p. 201–222.
- Eriksson, S.C., 1989, Phalaborwa: A saga of magmatism, metasomatism and miscibility, in *Bell, K., ed., Carbonatites: Genesis and evolution: London, Unwin Hyman*, p. 221–254.
- Finch, R.J., Hanchar, J.M., Hoskin, P.W.O., and Burns, P.C., 2000, Rare earth elements in synthetic zircon. 2. A single-crystal X-ray study of xenotime substitution: *American Mineralogist* (in press).
- Hanchar, J.M., and Hoskin, P.W.O., 1998, Mud Tank Carbonatite, Australia, zircon: *Society for Luminescence Microscopy and Spectroscopy Newsletter 10*, p. 2–3.
- Hanchar, J.M., Finch, R.J., Hoskin, P.W.O., Watson, E.B., Cherniak, D.J., and Mariano, A.N., 2000, Rare earth elements in synthetic zircon. 1. Synthesis, and rare earth element and phosphorus doping: *American Mineralogist* (in press).
- Heaman, L.M., Bowins, R., and Crocket, J., 1990, The chemical composition of igneous zircon suites: Implications for geochemical tracer studies: *Geochimica et Cosmochimica Acta*, v. 54, p. 1597–1607.
- Hinton, R.W., and Upton, B.G.J., 1991, The chemistry of zircon: Variations within and between large crystals from syenite and alkali basalt xenoliths: *Geochimica et Cosmochimica Acta*, v. 55, p. 3287–3302.
- Hoskin, P.W.O., 1998, Minor and trace element analysis of natural zircon (ZrSiO<sub>4</sub>) by SIMS and laser ablation ICPMS: A consideration and comparison of two broadly competitive techniques: *Journal of Trace and Microprobe Techniques*, v. 16, p. 301–326.
- Hoskin, P.W.O., and Black, L.P., 2000, Metamorphic zircon formation by solid-state recrystallisation of protolith igneous zircon: *Journal of Metamorphic Geology* (in press).
- Hoskin, P.W.O., Kinny, P.D., Wyborn, D., and Chappell, B.W., 2000, Identifying accessory mineral saturation during differentiation in granitoid magmas: An integrated approach: *Journal of Petrology* (in press).
- Ireland, T.R., and Wlotzka, F., 1992, The oldest zircons in the solar system: *Earth and Planetary Science Letters*, v. 109, p. 1–10.
- Ireland, T.R., Flöttmann, T., Fanning, C.M., Gibson, G.M., and Preiss, W.V., 1998, Development of the early Paleozoic Pacific margin of Gondwana from detrital-zircon ages across the Delamerian orogen: *Geology*, v. 26, p. 243–246.
- Maas, R., Kinny, P.D., Williams, I.S., Froude, D.O., and Compston, W., 1992, The Earth's oldest known crust: A geochronological and geochemical study of 3900–4200 Ma old detrital zircons from Mt. Narryer and Jack Hills, Western Australia: *Geochimica et Cosmochimica Acta*, v. 56, p. 1281–1300.
- Machado, N., Schrank, A., Noce, C.M., and Gauthier, G., 1996, Ages of detrital zircon from Archean-Paleoproterozoic sequences: Implications for Greenstone Belt setting and evolution of a Transamazonian foreland basin in Quadrilátero Ferrífero, southeast Brazil: *Earth and Planetary Science Letters*, v. 141, p. 259–276.
- McDonough, W.F., and Sun, S.-S., 1995, The composition of the Earth: *Chemical Geology*, v. 120, p. 223–253.
- Miller, J.M., and Gray, D.R., 1996, Structural signature of sediment accretion in a Palaeozoic accretionary complex, southeastern Australia: *Journal of Structural Geology*, v. 18, p. 1245–1258.
- Morton, A.C., 1992, Provenance of Brent Group sandstones: Heavy mineral constraints, in *Morton, A.C., et al., eds., Geology of the Brent Group: Geological Society [London] Special Publication 61*, p. 227–244.
- Morton, A.C., and Berge, C., 1995, Heavy mineral studies in the Statfjord and Nansen Formations of the Brent Field, North Sea: A new tool for reservoir subdivision and correlation: *Petroleum Geoscience*, v. 1, p. 355–364.
- Morton, A.C., Clauoué-Long, J.C., and Berge, C., 1996, SHRIMP constraints on sediment provenance and transport history in the Mesozoic Statfjord Formation, North Sea: *Geological Society of London Journal*, v. 153, p. 915–929.
- Murali, A.V., Parthasarathy, P., Mahadevan, T.M., and Sankar Das, M., 1983, Trace element characteristics, REE patterns and partition coefficients of zircons from different geological environments—A case study on Indian zircons: *Geochimica et Cosmochimica Acta*, v. 47, p. 2047–2052.
- Roback, R.C., and Walker, N.W., 1995, Provenance, detrital zircon U-Pb geochronometry, and tectonic significance of Permian to Lower Triassic sandstone in southeastern Quesnellia, British Columbia and Washington: *Geological Society of America Bulletin*, v. 107, p. 665–675.
- Sawka, W.N., 1988, REE and trace element variations in accessory minerals and hornblende from the strongly zoned McMurry Meadows pluton, California: *Royal Society of Edinburgh Transactions, Earth Sciences*, v. 79, p. 157–168.
- Schaltegger, U., Fanning, C.M., Günther, D., Maurin, J.C., Schulmann, K., and Gebauer, D., 1999, Growth, annealing and recrystallisation of zircon and preservation of monazite in high-grade metamorphism: Conventional and in situ U-Pb isotope, cathodoluminescence and microchemical evidence: *Contributions to Mineralogy and Petrology*, v. 134, p. 186–201.
- Sircombe, K.N., 1999, Tracing provenance through the isotope ages of littoral and sedimentary detrital zircon, eastern Australia: *Sedimentary Geology*, v. 124, p. 47–67.
- Speer, J.A., 1982, Zircon, in *Ribbe, P.H., ed., Orthosilicates: Mineralogical Society of America Reviews in Mineralogy*, v. 5, p. 67–112.
- Watson, E.B., 1996, Surface enrichment and trace-element uptake during crystal growth: *Geochimica et Cosmochimica Acta*, v. 60, p. 5013–5020.
- Watson, E.B., and Harrison, T.M., 1983, Zircon saturation revisited: Temperature and composition effects in a variety of crustal magma types: *Earth and Planetary Science Letters*, v. 64, p. 295–304.
- Williams, I.S., Buick, I.S., and Cartwright, I., 1996, An extended episode of early Mesoproterozoic fluid flow in the Reynolds Range, central Australia: *Journal of Metamorphic Geology*, v. 14, p. 29–47.
- Wyborn, D., Turner, B.S., and Chappell, B.W., 1987, The Boggy Plain Supersuite: A distinctive belt of I-type igneous rocks of potential economic significance in the Lachlan fold belt: *Australian Journal of Earth Sciences*, v. 34, p. 21–43.
- Young, D.N., and Black, L.P., 1991, U-Pb zircon dating of Proterozoic igneous charnockites from the Mawson Coast, East Antarctica: *Antarctic Science*, v. 3, p. 205–216.

Manuscript received December 15, 1999

Revised manuscript received March 24, 2000

Manuscript accepted April 4, 2000