

Chemical and isotopic study of thermal springs and gas discharges from Sierra de Chiapas, Mexico

A. Nencetti¹, F. Tassi¹, O. Vaselli^{1,2}, J. L. Macías³, G. Magro⁴, B. Capaccioni⁵, A. Minissale² and J.C. Mora³

¹ *Department of Earth Sciences, Florence, Italy*

² *CNR-Institute of Geosciences and Earth Resources, Florence, Italy*

³ *Institute of Geophysics, UNAM, Mexico D.F, Mexico*

⁴ *CNR-Institute of Geosciences and Earth Resources, Pisa, Italy*

⁵ *Institute of Volcanology and Geochemistry, Urbino, Italy*

Received: January 23, 2003; accepted: October 19, 2004

RESUMEN

La Sierra de Chiapas localizada en el Sureste de México, se caracteriza por la presencia de descargas de gas y agua. La mayoría de los manantiales termales se asocian a rocas volcánicas Terciarias a lo largo de fallas regionales con orientación NO-SE.

Las descargas termales se dividen en cuatro grupos: i) aguas con composición Na-Cl y Sólidos Disueltos Totales (SDT) >3000 mg/L; ii) aguas con composición Ca-SO₄ y valores de SDT entre 1400 y 2300 mg/L; iii) aguas con composición Na-Cl, bajos contenidos de SDT (800-2400 mg/L) y un contenido de sulfato alto (hasta 650 mg/L) y iv) aguas con una composición Ca-HCO₃ y salinidad baja (SDT <250mg/L). La mayoría de estas aguas están asociadas con descargas de gas compuestas por N₂ (hasta 93 % en vol.), CO₂ (2.4-31.2 % en vol.) y Ar (hasta 1.25 % en vol.) con origen predominantemente meteórico. H₂S aparece únicamente en las muestras de gas colectadas en El Azufre (hasta 1.1 % en vol.). Los valores de δ¹³C-CO₂ comúnmente inferiores a -9.7‰ (PDB) sugieren un origen parcialmente biogénico del CO₂.

Los parámetros químicos e isotópicos de los manantiales indican que la circulación de fluidos en la Sierra de Chiapas se regula principalmente por la infiltración de aguas meteóricas en rocas carbonatadas del Cretácico medio-tardío hasta las formaciones evaporíticas del Jurásico tardío-Cretácico temprano. Estas últimas representan la fuente principal de las especies en solución de las aguas.

No se encontraron evidencias de la presencia de sistemas con entalpías medias a altas por debajo de la Sierra de Chiapas.

PALABRAS CLAVE: Isótopos, química de manantiales, Chiapas, México.

ABSTRACT

Thermal water and gas discharges south-east of El Chichón volcano, Mexico are associated mainly with NW-SE oriented fault systems. Spring discharges include i) waters with Na-Cl composition and TDS >3000 mg/L; ii) waters with Ca-SO₄ composition and TDS values between 1400 and 2300 mg/L; iii) waters with Na-Cl composition and TDS of 800 to 2400 mg/L and sulphate content up to 650 mg/L and iv) waters with Ca-HCO₃ composition and low salinity (TDS <250mg/L). Most of these waters are associated with free-gas discharges of N₂ (up to 93 % by vol.), CO₂ (2.4 to 31.2 % by vol.) and Ar (up to 1.25 % by vol.) with a predominant meteoric origin. H₂S is present only in gas samples collected at El Azufre (up to 1.1 % by vol.). The δ¹³C CO₂ values are always below -9.7‰ (PDB) and suggest a partially biogenic origin for CO₂.

Chemical and isotopic features of spring discharges indicate that fluid circulation in the Sierra de Chiapas is mainly regulated by meteoric waters that tend to infiltrate the upper and middle-Cretaceous carbonate units up to the lower Cretaceous-upper Jurassic evaporitic formations (by López-Ramos, 1982). The latter provide the main source of the species in solution.

No evidence for high-to-medium enthalpy systems at depth beneath the Sierra de Chiapas has been found.

KEY WORDS: Isotopes, chemistry, thermal springs, water, Chiapas, Mexico.

1. INTRODUCTION

The Sierra de Chiapas is a NW-SE elongated area transected by regional faults (Meneses Rocha, 1991; García-Palomo *et al.*, 2004). It is located in the northern-central part of Chiapas State (Southeastern Mexico), where thermal water and gas discharges occur (Figure 1). They are often used as spas and for therapeutic purposes. Despite the economic interest of these manifestations, their chemical characteriza-

tion and origin are still poorly constrained. On the other hand, the thermal discharges located in the area of El Chichón volcano, after the violent eruption occurred on March 1982, have been extensively investigated (e.g. Taran *et al.*, 1998; Tassi *et al.*, 2003; Capaccioni *et al.*, 2004; Rouwet *et al.*, 2004).

This work presents the results of chemical and isotopic analyses of water samples of 9 thermal springs and 5 cold springs and gas discharges collected in November 1999. The

sites are located in a region that includes the cities of Pichucalco, San Cristobal de Las Casas, Venustiano Carranza and Tuxtla Gutiérrez in the Chiapanecan Volcanic Arc (CVA), where several Cenozoic volcanoes occur (Figure 1). The main goals of this paper are to: i) establish the origin of thermal spring discharges; ii) determine the chemical processes governing the composition and circulation of these fluids and iii) assess their possible relations with Cenozoic volcanism in the area.

2. TECTONICS AND GEOLOGICAL SETTING

The Sierra de Chiapas is a folded and faulted chain of Mesozoic and Tertiary sedimentary rocks located in the central and northern portions of Chiapas State (Figure 1). The fold axes trend north-west and generally plunge north-westwards beneath the Pliocene and younger sediments of the coastal Tabasco plain of the Gulf of Mexico, as observed in the northern part of El Chichón volcano (Duffield *et al.*, 1984; Macías *et al.*, 1997; García-Palomo *et al.*, 2004). Nixon (1982) proposed that an extensional tectonic regime was acting in the area probably related to the presence of a triple junction between the North American, Caribbean and Cocos plates. However, it is widely accepted that the volcanism in the region is related to the subduction of the Cocos plate beneath Mexico and Central America (Havskov *et al.* 1983; Burbach *et al.* 1984; Luhr *et al.*, 1984; Macías *et al.*, 2003). According to Damon and Montesinos (1978), El Chichón is the youngest volcano in the north-western part of the 150 km long CVA, a NW-SE volcanic belt within the Chiapas State. Capaul (1987) described the CVA as a Quaternary chain composed of low profile, small volume volcanoes (3 to 4 km³) that have erupted alkali-and-incompatible-rich andesitic rocks with respect to the Trans-Mexican Volcanic Belt and the Central America Volcanic Arc rocks.

The stratigraphic sequence of the sedimentary formations consists of: i) lower Cretaceous upper Jurassic evaporitic formations, which have locally formed salt diapirs, as resulted from drillings for petroleum explorations; ii) middle to upper Cretaceous formations of massive dolomite limestone; iii) Tertiary rocks that include Paleocene to Miocene formations, consisting predominantly of interbedded sandstones and siltstones (López-Ramos, 1979; De la Rosa *et al.*, 1989; Ferrusquia-Villafranca, 1996; and references therein, Figure 1).

With the exception of El Chichón volcano and Los Tuxtlas Volcanic Field, a gap in the distribution of the Holocene volcanoes extends 650 km from Tacaná, close to the Guatemala border, to Pico de Orizaba, at the eastern end of the Trans-Mexican Volcanic Belt. Differently, a large number of Cenozoic igneous rocks are present in Chiapas, pertaining to the CVA, and the Chiapas Massif (Damon and

Montesinos, 1978). The former is mainly composed by: i) the Santa Fe mining district-Selva Negra region, dated 2.79 Ma, consisting of volcanic and intrusive rocks and ii) the Cerro Tzontehuitz (2.14 Ma), a >3000 m high volcano close to San Cristobal de Las Casas; iii) the Cerro Lanza (0.85 Ma), a volcanic plug near Nicolás Ruíz characterized by conspicuous vertical banding and the appearance of a Peléan spine, iv) the Navenchauc volcano (0.43 Ma) west of San Cristobal de Las Casas, and the northernmost active Chichón volcano (0.21 Ma) (Damon and Montesinos, 1978). The Chiapas Massif consists of two intrusive belts: 1) a Miocene belt (15-29 Ma) composed of granodiorites, qz-monzonites, and tonalities, parallel to the Pacific coast to the vicinity of Motozintla village and 2) a Paleozoic belt composed of granite to granodiorite rocks, parallel and inland with respect to the Miocene batholith (Mugica, 1987) (Figure 1).

3. SAMPLING AND ANALYTICAL TECHNIQUES

All the water and gas discharges are located in proximity of the Cenozoic volcanic rocks belonging to the CVA (Figure 1). Temperature was measured with a H/g thermometer, pH and HCO₃⁻ by titration with HCl, and NH₄⁺ with the Nessler method. The SiO₂ contents in the water samples have been measured in the field (heteropoly blue method), whereas the remaining major and trace components were analyzed in the laboratory.

Gas samples, emerging as gentle small bubbles within thermal pools, were sampled using a plastic funnel up-side-down positioned and stored into two pre-evacuated 100cc thorion tapped tubes, one containing 20 ml of a 4M NaOH solution (Giggenbach, 1975) to concentrate the non-reactive gas compounds in the head-space of the tubes. Gas chromatographic analyses for CO₂, H₂S, N₂, H₂, He, O₂, Ar and Ne contents were measured by a Shimadzu 15A gas-chromatograph equipped with a TCD (Thermal Conductivity Detector), while CO, after its conversion to CH₄ at 400°C, and hydrocarbon compounds were analyzed with a Shimadzu 14A gas-chromatograph equipped with a FID (Flame Ionization Detector). Analytical precision was <1% for major components and <5% for minor and trace compounds.

δ¹⁸O and δ²H values in liquid phase were determined with a Finnigan 250 Delta mass spectrometer after equilibrating the waters with CO₂ and after the reaction of 10 μ of water with 0.3 g of pure Zn at 500°C, respectively. The analytical errors were <10% for the abundance of elements and a precision of ±0.15 and ±1.1 ‰ for oxygen and hydrogen isotopes, respectively. The ¹³C/¹²C ratios of CO₂ in the gas phases, were determined using a Finnigan 250 Delta mass spectrometer. Several internationally accepted standards were used, with an error of ±0.05‰ (PDB). ³He/⁴He (expressed as R/Ra, where R is the measured ratio and Ra is that in the air,




Fig. 1. A Sketch map of southern Mexico showing the location of the Chiapanecan Volcanic Arc (CVA), the Central America Volcanic Arc (CAVA), and Los Tuxtlas Volcanic Field (TVF). B Regional map of the Sierra de Chiapas region (modified after López-Ramos, 1979) and location of the water and gas discharge sampling sites. Note the location of springs and their vicinity to the Cenozoic volcanic centers of the CVA.

1.39×10^{-6} ; Mamyrin and Tolstikhin, 1984) ratios in the gas phase were determined with a Map 215-50 magnetic mass spectrometer, equipped with an ion counting device. Resolution was close to 600 AMU for HD- ^3He at 5% of the peak. The He/Ne ratio is measured with a quadrupole mass spectrometer (VG-Spectralab 200). Both the spectrometers are connected to a metal high-vacuum line, where gas samples are processed to separate noble gases from the gaseous matrix. The reproducibility of R/Ra and He/Ne ratios in air standard is better than 5 % (Minissale et al., 2000; Magro et al., 2003).

4. CHEMICAL COMPOSITION OF THE FLUID PHASES

4.1. Water chemistry

Water samples have temperatures ranging between 11°C and 38.5 °C and pH varying from slightly acidic (5.77) to slightly alkaline (7.61). The analytical results for the main components and some trace compounds are listed in Table 1. On the basis of both Total Dissolved Solids (TDS, Table 1) and relative abundance of HCO_3^- , SO_4^{2-} and Cl^- ions (Figure 2), the Sierra de Chiapas waters can be divided, as follows: i) “Group I” (samples: Poza La Virgen, El Carmen I and II), that plots in the field pertaining to Cl^- , with a Na-Cl composition and TDS > 3000 mg/L; ii) “Group II” (samples: Villa Las Rosas and El Paraiso), positioned close to the SO_4^{2-} corner, with a Ca- SO_4 composition and TDS values between 1400 and 2300 mg/L; iii) “Group III” (samples: El Azufre Piscina I and II, El Azufre lake and El Carmen fria) with a

Na-Cl composition and differing from “Group I” for having a lower salinity (TDS ranging from 800 to 2400 mg/L) and a higher sulfate content (up to 650 mg/L); iv) “Group IV” (samples: El Azufre fría, Villa Las Rosas fría, Cerro Tzontehuitz, Chimiljotepec, Carranza Manantial), close to the HCO_3^- corner, characterized by a Ca- HCO_3 composition and low salinity (TDS < 250 mg/L).

The strong positive correlation between Na^+ and Cl^- (reported as meq/L) contents (Figure 3) in the Sierra de Chiapas waters suggests a common origin for these two ions and, possibly, resulting from the interaction with evaporitic material (dated lower Cretaceous-upper Jurassic by López-Ramos, 1979). As shown in Figure 4, with the only exception of waters pertaining to “Group IV”, the $(\text{Ca}^{2+} + \text{Mg}^{2+}) / (\text{HCO}_3^- + \text{SO}_4^{2-})$ ratios (as meq/L) are close to 1, again suggesting that the contents of these chemical compounds are mainly related to dissolution processes of evaporitic material. NH_4^+ and H_3BO_3 contents show large variations (from 0.09 to 22 mg/L and from <0.01 and 16 mg/L, respectively) and generally display a good correlation with Cl^- contents and TDS (Table 1).

Nitrate contents vary between 0.11 and 4.5 mg/L, with the exception of the Carranza spring, where anthropic pollution cannot be excluded. Bromide values (ranging from 0.01 to 2.2 mg/L) show a positive correlation with Cl^- contents (Figure 5). F⁻ abundance goes from 0.02 to 1.8 mg/L while those of Li^+ vary from 0.02 to 0.66 mg/L. SiO_2 contents are between 11 to 31 mg/L, showing clear correlation with the outlet temperatures. Silica concentrations of El Azufre springs, determined by using the heteropoly blue method,

Table 1

Chemical composition of mineral and cold waters of Sierra de Chiapas

Tippe	Date	T°C	pH	TDS	Na ⁺	K ⁺	Mg ²⁺	Ca ²⁺	NH ₄ ⁺	Li ⁺	SiO ₂	HCO ₃ ⁻	Cl ⁻	SO ₄ ²⁻	F ⁻	Br ⁻	NO ₃ ⁻	H ₃ BO ₃	δ ¹⁸ O	δ ² H	
El Azufre P 1	tsg	nov 99	31.5	6.28	2408	478	29	90	274	22	0.34	n.a.	720	842	621	1.0	1.9	0.11	4.98	-4.5	-21.0
El Azufre P 2	tsg	nov 99	31.3	6.5	2410	456	28	95	338	21	0.33	n.a.	701	760	650	1.1	1.8	0.12	6.29	-4.3	20.0
El Azufre Natural	ts	nov 99	29	6.57	1573	284	19	62	230	15	0.22	n.a.	537	445	460	1.0	1.2	0.18	3.20	n.a.	n.a.
El Azufre fría	s	nov 99	23	7.56	148	5.4	1.2	19	53	0.08	<0.001	n.a.	207	8.5	28	0.060	0.025	2.9	1.09	-4.6	-21.4
Tzontehuitz Volcano	s	nov 99	11	6.2	23	1.4	0.71	0.34	2.1	0.19	<0.001	16	9.1	0.6	1.8	0.02	<0.01	<	<0.01	n.a.	n.a.
Chimiljotepec	s	nov 99	14	5.77	30	3.2	1.9	0.90	2.8	0.2	<0.001	19	21	0.5	1.0	0.06	0.01	0.47	<0.01	n.a.	n.a.
Bal. Villa Las Rosas	tsg	nov 99	23.4	6.86	2286	1.7	2.0	131	525	3.4	0.04	18	287	2.7	1600	1.8	0.025	0.11	<0.01	-9.8	-64.0
Bal. V. Las Rosas fria	s	nov 99	22	7.03	161	1.9	1.1	26	112	1.9	<0.001	11	464	1.4	0.9	0.05	<0.01	4.5	<0.01	-8.8	-54.6
Bal. El Paraiso	ts	nov 99	25	7.51	1438	15	1.7	70	343	1.4	0.02	11	271	28	960	0.36	0.05	7.6	<0.01	-9.3	-65.5
Carranza Manantial	s	nov 99	27	7.61	280	10	0.5	141	19	1.9	<0.001	19	720	2.1	18	0.08	<0.01	68	3.03	n.a.	n.a.
Bal. El Carmen I	tsg	nov 99	38.5	6.46	3092	855	72	61	255	1.6	0.66	24	744	1330	490	0.61	1.6	0.30	<0.01	-8.0	-53.7
Bal El Carmen II	tsg	nov 99	38.5	6.55	3013	790	73	65	250	2.2	0.64	21	653	1300	505	0.68	2.2	1.8	5.43	-8.3	-55.3
Bal. El Carmen fría	ts	nov 99	30	6.7	874	180	17	32	141	3.6	0.15	20	367	265	210	0.18	0.40	4.2	<0.01	-8.3	-53.6
Poza La Virgen	tsg	nov 99	37	6.44	3261	900	79	64	259	1.9	0.14	31	708	1410	510	0.51	1.5	0.33	16.01	-8.4	-56.3

All the species are expressed in mg/L; TDS expressed in mg/L; n.a.=not analysed; tsg=thermal spring with gas associated; ts=thermal spring; s=cold spring.




Fig. 2. $\text{SO}_4^{2-}\text{-Cl-HCO}_3^-$ ternary diagram for discharges from Sierra de Chiapas region. Open squares are cold springs ($T < 23^\circ\text{C}$); closed circles are: thermal springs ($T > 23^\circ\text{C}$).




Fig. 3. Na^+ versus Cl^- (in meq/L) diagram for water discharges from Sierra de Chiapas region. Ion contents are in mg/L. Symbols as in Fig. 2.

were up to 60 mg/L. However, these values are jeopardized by the high H_2S contents (Table 2) and are not considered in the present work.

The $\text{Na}/400$, $\text{K}/100$ and $\sqrt{\text{Mg}}$ triangular diagram (Figure 6; Giggenbach, 1988) shows that all water samples fall within the immature water field and aligned along the water-rock dissolution line; thus, K/Na and K^2/Mg geothermometers can not be applied to obtain a real assessment of the reservoir equilibrium temperature. The silica geothermometers




Fig. 4. $\text{Ca}^{2+} + \text{Mg}^{2+}$ vs. $\text{HCO}_3^- + \text{SO}_4^{2-}$ diagram (in meq/L) for the water discharges from Sierra de Chiapas region. Symbols as in Fig. 2.




Fig. 5. Br^- vs. Cl^- diagram (in meq/L) for the water discharges from Sierra de Chiapas region.

used (Fournier and Rowe, 1966; Giggenbach and Corrales, 1988), applied to all the thermal waters but El Azufre, present more homogeneous values and range from 40°C to 80°C , although these temperatures have to be regarded as minimum equilibrium temperatures due to the mixing processes between deep and superficial waters.

$\delta^{18}\text{O}$ values vary between -4.3‰ at El Azufre Piscina

II and - 9.8 ‰ at Villa Las Rosas, whereas $\delta^2\text{H}$ values range from -20.0 ‰ SMOW at El Azufre Piscina II and -65.5 ‰ at El Paraiso (Table 1). As shown in the binary diagram of Figure 7, the $\delta^{18}\text{O}$ - $\delta^2\text{H}$ values for all samples investigated approach the Global Meteoric Water Line (GMWL, Craig, 1963), suggesting that all the water discharges have a meteoric origin.

4.2. Gas chemistry

The chemical composition (in % by vol.) of gas samples is reported in Table 2. N_2 is always the main component (up to 93 %), while CO_2 content varies from 2.4 % to 31.2 %. O_2 contents are very low or under the detection limit (0.001 %), while high contents of Ar, larger than the concentration in air (0.93 %), have been measured in all samples (up to 1.25%). H_2S is present in detectable amounts (>0.005 %) only in the gas samples collected at El Azufre (up to 1.1 %).

Among hydrocarbons, CH_4 is the most abundant component (up to 5.0 % in vol.), but significant amounts of ethane, propane, *i*-butane, *1*-butene and *i*-butene have also been detected (up to 1.2 % of the total organic compound contents) (Table 2). He contents are of the order of some hundreds of ppm by vol., while H_2 and Ne contents are from 2 to 20 and from 5.4 to 9.5 ppm by vol., respectively. CO is always below the instrumental detection limit (<0.1 ppm by vol).

In the $10\text{He}-\text{N}_2/100-\text{Ar}$ ternary diagram (Figure 8) all gas samples plot in the field defined by air, ASW (Air-Saturated Water, Giggenbach *et al.*, 1983) and the He corner, indicating that N_2 and Ar are mostly atmospheric-originated, while considerable amounts of non-atmospheric He (with values up to 363 ppmv, Table 2) were added to the circulating fluids.

R/Ra values (Table 2) range between 0.6 at El Azufre




Fig. 6. Triangular diagram of Na/400-K/10-√Mg (Giggenbach and Corrales, 1992).

Piscina I sample and 1.0 at El Carmen sample, which may indicate, if we consider a value of $R/R_a = 8$ for the a MORB-type mantle (Martel *et al.*, 1989; Ozima *et al.*, 2002), a contribution of 7-13% of mantle-derived ^3He . The $\delta^{13}\text{C}$ in CO_2 values (Table 2) range between -9.7 ‰ to -12.5 ‰ PDB. These isotopic values seem to be in relation with decomposition of organic material (up to $\delta^{13}\text{C} = -20$ ‰ PDB), although mixing with a deeper-originated (mantle- or thermometamorphic-related) end-member (possibly characterized by $-7 < \delta^{13}\text{C} < +1$) cannot be excluded.

5. DISCUSSION AND CONCLUSIONS

The chemical and isotopic data of sampled water and gas discharges in the Sierra de Chiapas, Mexico may give some insights on the origin and on the chemical evolution of

Table 2

Chemical composition of gas discharges from Sierra de Chiapas

	Date	CO_2	N_2	H_2S	Ar	O_2	CH_4	He	H_2	Ne	CO	Ethane	Propane	<i>i</i> -Butane	<i>1</i> -Butene	<i>i</i> -Butene	$\delta^{13}\text{C}-\text{CO}_2$	R/R _a	
El Azufre P 1	nov 99	18.24	79.0	0.82	1.10	0.005	1.10	162	13.20	8.7	<0.01	83	1.2	0.27	0.18	0.09	-10.5	0.57	
El Azufre P 1a	nov 99	18.21	79.0	0.86	1.03	0.005	1.11	263	15.60	7.8	<0.01	133	2.0	0.62	0.09	0.31	-9.7		
El Azufre P 2	nov 99	29.25	68.0	0.55	0.99	0.014	0.95	208	53.00	7.7	<0.01	115	1.6	0.40	0.04	0.08	-12.3		
El Azufre lake	nov 99	2.39	93.0	1.10	1.21	0.330	1.59	274	8.66	9.4	<0.01	156	2.3	0.73	0.05	0.11	-11.3		
Bal. El Carmen	nov 99	30.70	63.0	<0.01	0.85	0.570	5.04	363	5.42	5.7	<0.01	5.8					9.7	1.03	
Poza la Virgen	nov 99	31.17	53.0	<0.01	0.74	11.000	4.57	249	20.70	5.4	<0.01	14				0.92	9.7	1.03	
Bal Villa																			
Las Rosas	nov 99	3.01	95.0	<0.01	1.25	0.730	0.03	15.5	2.00	9.5	<0.01	1.0				0.11	12.5	0.66	

All data are in % by volume, with the exception of He, H_2 , Ne, Co, Etane, Propane, *i* Butane, *1*-Butane, *i*-Butene that are in ppmv.




Fig. 7. $\delta^{18}\text{O}$ - $\delta^2\text{H}$ diagram for water discharges from Sierra de Chiapas region. The Global Meteoric Water Line (after Craig, 1963) is reported.




Fig. 8. $\text{N}_2/100$ -Ar-He*10 ternary diagram for gas discharges from Sierra de Chiapas region (after Giggenbach, 1993).

the thermal fluids in the region. The TDS values are plotted versus the percentage of HCO_3^- with respect to the sum of total anions ($\text{HCO}_3^- + \text{SO}_4^{2-} + \text{Cl}^-$) and reported in Figure 9. It can be speculated that water chemistry may be related to mixing processes between a superficial aquifer, represented by the cold spring discharges ("Group IV" samples), and a more saline end member, characterized by a composition similar to that of both Poza La Virgen and Balneario El Carmen I and II waters ("Group I" samples). The striking

correspondence between the location of the thermal springs and the outcrops of the Cenozoic volcanic rocks (Figure 1) suggests that fluid circulation is controlled by the main fracture and fault systems (Nencetti 2001).

The predominance of atmospheric-related components in all gas phases sampled (Table 2) and the $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values of all spring samples (Table 1; Figure 5) suggests that the presence of active high enthalpy hydrothermal systems at relatively shallow depths seems unlikely. This is in accordance with recent investigations on the geothermal resources of Chiapas (Prol Ledesma and Juárez, 1986), which indicate deep fluid temperatures ranging from 85°C to 110°C . Chemical and isotopic considerations may allow to hypothesize a possible scheme for the evolution of the circulating fluids. Meteoric waters easily infiltrate through the south-east oriented outcrops of upper and middle-Cretaceous carbonate units, which extend from Tuxtla Gutiérrez and Venustiano Carranza up to Pichucalco (Figure 1). The permeability of these rocks is possibly enhanced by the fault systems related to the regional tectonics (López-Ramos, 1979; Canul and Rocha, 1982; Nixon, 1982; Macías *et al.*, 1997; Meneses Rocha, 1991; García-Palomo *et al.*, 2004). The typical atmospheric gases such as N_2 , O_2 , Ar and Ne enrich the groundwater during the infiltration process, as dissolved gas species in the rainfall water. Therefore, permeated meteoric waters reach the evaporitic rocks (*i.e.* anhydrite halite-carbonatic formation) that are located at variable depth (Montes de Oca, 1979; López Ramos, 1979; Duffield *et al.*, 1984). During the circulation, prolonged water-rock interactions occur and the temperature of the waters tend to increase as they deepen. Contemporaneously, CO_2 , possibly also deriving from deep sources (mantle- and/or thermometamorphic-derived), and He deriving from radiogenic decay in the crustal environment, are produced and added to these thermal fluids. However, as indicated by R/Ra values of Poza La Virgen and El Carmen II gas discharges, which emergence locations are in coincidence with the presence of igneous bodies of upper Pleistocene age (Figure 1), some contribution of mantle ^3He , related to the active tectonic regime of the area, cannot be excluded. Moreover, at shallow level, during the ascending phase of the convective circuit, some isotopically light CO_2 , derived from the alteration of organic matter (as supported by the $\delta^{13}\text{C}$ values), H_2S and hydrocarbon compounds further modify the chemical characteristics of the gas thermal discharges. In conclusion, the chemical and isotopic characteristics of the Sierra de Chiapas gas and water discharges, suggest that the presence of active high enthalpy hydrothermal systems at relatively shallow depths seems to be unlikely.

ACKNOWLEDGEMENTS

This study was supported by a CNR-CONACYT Bilateral Project and by the Italian Space Agency (ASI, Resp. OV). Many thanks are due to M. Paolieri for his assistance




Fig. 9. TDS (in ppm) and HCO₃⁻ (in % with respect to the sum of HCO₃⁻, SO₄²⁻ and Cl⁻) for the water discharges from Sierra de Chiapas region. Symbols as in Fig. 2.

during the stable isotopic analysis. Yuri Taran and an anonymous reviewer are warmly thanked for their criticism and suggestions.

BIBLIOGRAPHY

BURBACH, V. G., C. FROHLICH, D. W. PENNINGTON and T. MATUMOTO, 1984. Seismicity and tectonics of the subducted Cocos Plate. *J. Geophys. Res.*, 89, 7719-7735.

CANUL, R. F. and V. L. ROCHA, 1982. Informe geológico de la zona geotérmica de "El Chichónal", Chiapas. Unpublished Report Comisión Federal De Electricidad, Morelia, Michoacán, México, 38p.

CAPAU, W. A., 1987. Volcanoes of the Chiapas volcanic belt, Mexico. Thesis. Michigan Technological University, pp 93.

CAPACCIONI, B., Y. TARAN, F. TASSI, O. VASELLI, F. MANGANI and J. L. MACÍAS, 2004. Source conditions and degradation processes of light hydrocarbons

in volcanic gases: an example from El Chichón volcano (Chiapas State, Mexico). *Chem. Geol.*, 206, 81-96.

CRAIG, H., 1963. Isotopic variations in meteoric waters. *Science*, 123, 1702-1703.

DAMON, P. E. and E. MONTESINOS, 1978. Late Cenozoic volcanism and metallogenesis over an active Benioff zone in Chiapas, Mexico. *Arizona Geol. Soc. Digest*, 11, 155-168.

DE LA ROSA, J. L., A. EBOLI and M. DÁVILA, 1989. Geología del Estado de Chiapas. Subdirección de Construcción, Comisión Federal de Electricidad. Harla, Mexico City, 192 p.

DUFFIELD, W. A., R. I. TILLING and R. CANUL, 1984. Geology of El Chichón Volcano, Chiapas, Mexico. *J. Volcanol. Geotherm. Res.*, 20, 117-132.

FERRUSQUIA-VILAFRANCA, I., 1996. Contribución al conocimiento geológico de Chiapas-El área de Ixtapa-

- Soyaló. Boletín 109, Instituto de Geología, UNAM, 130 p.
- FOURNIER, R. O. and J. J. ROWE, 1966. Estimation of underground temperatures from the silica content of water from hot spring and wet-steam wells. *Am. Jour. Scie.*, 264, 685-697.
- GARCÍA-PALOMO, A., J. L. MACÍAS and J. M. ESPÍNDOLA, 2004. Strike-slip faults and K-Alkaline volcanism at El Chichón volcano, southeastern Mexico. *J. Volcanol. Geotherm. Res.*, 136, 247-268.
- GIGGENBACH, W. F., 1975. A simple method for the collection and analyses of volcanic gas sample. *Bull. Volcanol.*, 39, 132-145.
- GIGGENBACH, W. F., 1988. Geothermal solute equilibria. Derivation of Na-K-Mg-Ca geoindicators. *Geoch. Cosmoch. Acta*, 52, 2749-2765.
- GIGGENBACH, W. F. and R. CORRALES, 1992. Isotopic and chemical composition of water and steam discharges from volcanic-magmatic-hydrothermal system of the Guanacaste Geothermal Province, Costa Rica. *Appl. Geochem.*, 7, 309-322.
- GIGGENBACH, W. F., R. GONFIANTINI, B. L. JANGI and A. H. TRUESDELL, 1983. Isotopic and chemical composition of Parbaty valley geothermal discharges, NW-Himalaya. *Geotherm.*, 12, 199-222.
- HAVSKOV, J., S. DE LA CRUZ-REYNA, S. SINGH, F. MEDINA and C. GUTIÉRREZ, 1983. Seismic activity related to the March-April, 1982 eruptions of El Chichón volcano, Chiapas, Mexico. *Geophys. Res. Lett.*, 10, 293-296.
- LÓPEZ-RAMOS, E. L., 1979. Geología de México. Edición Escolar, Tomo III, segunda edición, México, D.F., México, 446 pp.
- LUHR, J. F., I. S. E. CARMICHAEL and J. C. VAREKAMP, 1984. The 1982 eruptions of El Chichón Volcano, Chiapas, Mexico: mineralogy and petrology of the anhydrite-bearing pumices. *J. Volcanol. Geotherm. Res.*, 23, 69-108.
- MACÍAS, J. L., J. M. ESPÍNDOLA, Y. TARAN and P. A. GARCÍA, 1997. Explosive volcanic activity during the last 3,500 years at el Chichon volcano, Mexico. IAVCEI, General Assembly, Puerto Vallarta, Mexico. Field trip guide, 53 p.
- MACÍAS, J. L., J. L. ARCE, J. C. MORA, J. M. ESPÍNDOLA, R. SAUCEDO and P. MANETTI, 2003. The ~550 BP plinian eruption of el Chichon volcano, Chiapas, Mexico: Explosive volcanism linked to reheating of a magma chamber. *J. Geophys. Res.*, 108(b12), 2569.
- MAGRO, G., G. RUGGIERI, G. GIANELLI, S. BELLANI and G. SCANDIFFIO, 2003. Helium isotopes in paleofluids and present-day fluids of the Larderello geothermal field: Constraints on the heat source. *J. Geophys. Res.*, 108(B9), doi: 10.1029-/2001JB001590, 2003.
- MARTEL, D. J., J. DEAK, P. DOVENYI, F. HORVATH, R. K. O'NIONS, E. R. OXBURGH, L. STEGENA and M. STUTE, 1989. Leakage of helium from pannonian basin. *Nature*, 342, 908-912.
- MAMYRIN, B. A. and I. N. TOLSTIKHIN, 1984. Helium isotopes in natura. In: Fyfe, W.S. (Ed), *Developments in Geochemistry*, Series 3, Elsevier, Amsterdam.
- MENESES-ROCHA, J. J., 1991. Tectonic Development of the Ixtapa Graben, Chiapas, Mexico. Ph. D., University of Texas, Austin, 308 pp.
- MINISSALE, A., G. MAGRO, G. MARTINELLI, O. VASELLI and F. TASSI, 2000. Fluid geochemical transect in the Northern apennines(central-northern Italy): fluid genesis and migration and tectonic implications. *Tectonophysics*, 319, 199-222.
- MONTES DE OCA, S., 1979. Geología petrolera de la Sierra de Chiapas. *Asoc. Mex. Geol. Pétrol., Bol.*, 31 (1-2), 67-97.
- MUGICA, R., 1987, Estudio petrogenético de las rocas ígneas y metamórficas en el Macizo de Chiapas: Instituto Mexicano del Petróleo, C-2009.
- NENCETTI, A., 2001. Geochimica dei fluidi del Vulcano El Chichón (Chiapas, Messico). Unpublished Ms Thesis, University of Florence (Italy), pp 181. (In Italian).
- NIXON, G. T., 1982. The relationship between Quaternary volcanism and the seismic structure of subducted ocean lithosphere. *Bull. Geoc. Soc. Am.*, 93, 514-523.
- OZIMA, M. and F. A. PODOSEK, 2002. Noble gas geochemistry. Cambridge University Press, Second Edition, 286 pp.

- PROL-LEDESMA, R. M. and M. G. JUÁREZ, 1986. Geothermal map of Mexico. *J. Volcanol. Geotherm. Res.*, 28, 351-362.
- ROUWET, D., Y. A. TARAN and N. R. VARLEY, 2004. Dynamics and mass balance of El Chichón crater lake, Mexico. *Geoffs. Int.*, 43, 427-434.
- TARAN, Y., T. P. FISCHER, B. POKROVSKY, Y. SANO, M. A. ARMIENTA and J. L. MACÍAS, 1998. Geochemistry of the volcano-hydrothermal system of El Chichón Volcano, Chiapas, Mexico. *Bull. Volcanol.*, 59, 436-449.
- TASSI, F., O. VASELLI, B. CAPACCIONI, J. L. MACÍAS, A. NENCETTI, G. MONTEGROSSI and G. MAGRO, 2003. Chemical composition of fumarolic gas and spring discharges from El Chichón volcano, Mexico: causes and implications of the changes detected over the period 1998-2000. *J. Volcanol. Geotherm. Res.*, 123, 105-121.
- XU, S., S. NAKAI, H. WAKITA, X. WANG and X. FENG, 1997. Effects of hydrothermal processes on the chemical and isotopic composition of mantle-derived gases in SE China. Pergamon, *Geothermics*, 26, 2, 179-192.
-
- A. Nencetti¹, F. Tassi¹, O. Vaselli^{1,2,*}, J. L., Macías³, G. Magro⁴, B. Capaccioni⁵, A. Minissale², J. C. Mora³
- ¹ Department of Earth Sciences, Via G. La Pira 4, 50121 Florence, Italy
² CNR-Institute of Geosciences & Earth Resources, Via G. La Pira 4, 50121 Florence (Italy)
³ Institute of Geophysics, UNAM, D.F., 04510 Mexico City (Mexico)
⁴ CNR-Institute of Geosciences & Earth Resources, Via G. Moruzzi 1, 56124 Pisa (Italy)
⁵ Institute of Volcanology and Geochemistry, Loc. La Crocicchia, Ex-Sogesta 61029 Urbino (Italy)
- * Corresponding author: Orlando Vaselli, Department of Earth Sciences, Via G. La Pira, 4 - 50121 Florence (Italy).
Tel: +39 055 2756289; Fax: +39 055 284571;
Email: orlando@geo.unifi.it